

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
17 June 2004 (17.06.2004)

PCT

(10) International Publication Number
WO 2004/050094 A1

(51) International Patent Classification⁷: **A61K 31/495**,
A61P 37/00

(21) International Application Number:
PCT/EP2003/005491

(22) International Filing Date: 26 May 2003 (26.05.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2080064.5 3 December 2002 (03.12.2002) EP

(71) Applicant (*for all designated States except US*): **UCB, S.A.**
[BE/BE]; Allée de la Recherche 60, B-1070 Brussels (BE).

(72) Inventor; and

(75) Inventor/Applicant (*for US only*): **REVIRRON,**
Christophe [FR/CH]; Rossistrasse 25, CH-1735 Giffers
(CH).

(74) Agent: **LECHIEN, Monique**; UCB, S.A., Intellectual
Property Dept., Allée de la Recherche 60, B-1070 Brussels
(BE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO,
SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: USE OF LEVOCETIRIZINE FOR THE TREATMENT OF PERSISTENT ALLERGIC RHINITIS

(57) Abstract: The present invention relates to a pharmaceutical use of levocetirizine for the treatment of persistent allergic rhinitis.

WO 2004/050094 A1

USE OF LEVOCETIRIZINE FOR THE TREATMENT OF PERSISTENT ALLERGIC RHINITIS

The present invention relates to the use of levocetirizine for the preparation of drugs effective for the treatment of the persistent allergic rhinitis.

5 International patent application 94/06429 describes a method utilising levocetirizine for the treatment of seasonal and perennial allergic rhinitis.

It has now surprisingly been found that levocetirizine possesses therapeutic properties which render it particularly useful in the treatment of persistent allergic rhinitis. These activities are not observed in the dextrocetirizine.

10 The purpose of the invention concerns the treatment of persistent allergic rhinitis.

The present invention is based on the unexpected recognition that administration of pharmaceutical compositions comprising levocetirizine, or a pharmaceutically acceptable salt thereof to a patient treats the persistent allergic
15 rhinitis.

The present invention encompasses a method for treating persistent allergic rhinitis which comprises administering to a patient a therapeutically effective amount of levocetirizine or a pharmaceutically acceptable salt thereof.

The present invention also encompasses the use of levocetirizine or a
20 pharmaceutically acceptable salt thereof for the preparation of a medicament intended for the treatment of persistent allergic rhinitis.

The present invention relates to the use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for decreasing the symptoms of persistent allergic rhinitis and improving the quality of life.

25 In another aspect, the present invention relates to a method of treating in a patient persistent allergic rhinitis by administering an effective dose of levocetirizine or a pharmaceutically acceptable salt thereof.

The term "cetirizine" refers to the racemate of [2-[4-[(4
chlorophenyl)phenylmethyl] -1-piperazinyl]ethoxy]-acetic acid and its dihydrochloride
30 salt which is well known as cetirizine dihydrochloride; its levorotatory and dextrorotatory enantiomers are known as levocetirizine and dextrocetirizine. Processes for preparing cetirizine, an individual optical isomer thereof or a pharmaceutically acceptable salt thereof have been described in European Patent 0 058 146, Great Britain Patent 2.225.320, Great Britain Patent 2.225.321, United States Patent
35 5,478,941, European Patent application 0 601 028, European Patent Application 0 801 064 and International Patent Application WO 97/37982.

The term "levocetirizine" as used herein means the levorotatory enantiomer of cetirizine. More precisely, it means that the active substance comprises at least 90%

by weight, preferably at least 95% by weight, of one individual optical isomer of cetirizine and at most 10% by weight, preferably at most 5% by weight, of the other individual optical isomer of cetirizine. Each individual optical isomer may be obtained by conventional means, i.e., resolution from the corresponding racemic mixture or by asymmetric synthesis. Each individual optical isomer may be obtained from its racemic mixture by using conventional means such as disclosed in British patent application No. 2,225,321. Additionally, each individual optical isomer can be prepared from the racemic mixture by enzymatic biocatalytic resolution, such as disclosed in U.S. Patents No. 4,800,162 and 5,057,427.

10 The term "pharmaceutically acceptable salts" as used herein refers not only to addition salts with pharmaceutically acceptable non-toxic organic and inorganic acids, such as acetic, citric, maleic, succinic, ascorbic, hydrochloric, hydrobromic, sulfuric, and phosphoric acids and the like, but also its metal salts (for example sodium or potassium salts) or ammonium salts, the amine salts and the aminoacid salts. The best results have been obtained with levocetirizine dihydrochloride.

By patient, we understand children, adolescents and adults.

By the term "allergic rhinitis", we understand a symptomatic disorder of the nose induced by an IgE-mediated inflammation after allergen exposure of the membrane of the nose. Symptoms of allergic rhinitis include rhinorrhea, nasal obstruction, nasal itching, sneezing, ocular pruritis. The term "persistent allergic rhinitis", as used herein, refers to a disease when symptoms last more than 4 days per week and for more than 4 weeks. It is subdivided into mild and moderate-severe rhinitis. It is said "mild" when there are normal sleep, or no impairment of normal daily activities, sport, leisure, normal work and school, or no troublesome symptoms. It is said "moderate-severe" when there are abnormal sleep, or impairment of daily activities, sport, leisure, or problems caused at work or school, or troublesome symptoms.

A therapeutically effective amount of levocetirizine or a pharmaceutically acceptable salt thereof is used to treat or alleviate the effects of persistent allergic rhinitis. The dosage depends essentially on the specific method of administration and on the purpose of the treatment. The size of the individual doses and the administration program can best be determined based on an individual assessment of the relevant case. The methods required to determine the relevant factors are familiar to the expert.

35 A preferred daily dosage provides from about 0,0005 mg to about 2 mg of levocetirizine or a pharmaceutically acceptable salt thereof, per kg of body weight per patient. A particularly preferred daily dosage is from about 0,001 to about 2 mg per kg of body weight per patient. The best results have been obtained with a daily dosage

from about 0,005 to 1 mg per kg of body weight per patient. The dosage may be administered once per day of treatment, or divided into smaller dosages, for examples 1 to 4 times a day, and preferably 1 to 3 times a day, and administrated over about a 24 hours time period to reach a total given dosage. Bests results have been obtained with an administration of a compositions of the invention are twice a day for children; and 5 mg once a day for adults. The exact dosages in which the compositions are administered can vary according to the type of use, the mode of use, the requirements of the patient, as determined by a skilled practitioner. The exact dosage for a patient may be specifically adapted by a skilled person in view of the severity of the condition, the specific formulation used, and other drugs which may be involved.

Pharmaceutical compositions used according to the present invention may be administered by any conventional means. The routes of administration include intradermal, transdermal, slow release administration, intramuscular, oral and intranasal routes. Any other convenient route of administration can be used, for example absorption through epithelial or mucocutaneous linings.

The pharmaceutical forms according to the present invention may be prepared according to conventional methods used by pharmacists. The forms can be administered together with other components or biologically active agents, pharmaceutically acceptable surfactants, excipients, carriers, diluents and vehicles.

The pharmaceutical compositions of the invention include any conventional therapeutical inert carrier. The pharmaceutical compositions can contain inert as well as pharmacodynamically active additives. Liquid compositions can for example take the form of a sterile solution which is miscible with water. Furthermore, substances conventionally used as preserving, stabilizing, moisture-retaining, and emulsifying agents as well as substances such as salts for varying the osmotic pressure, substances for varying pH such as buffers, and other additives can also be present. If desired an antioxidant can be included in the pharmaceutical compositions.

Pharmaceutical acceptable excipients or carriers for compositions include saline, buffered saline, dextrose or water. Compositions may also comprise specific stabilizing agents such as sugars, including mannose and mannitol. Carrier substances and diluents can be organic or inorganic substances, for example water, gelatine, lactose, starch, magnesium stearate, talc, gum arabic, polyalkylene glycol and the like. A prerequisite is that all adjuvants and substances used in the manufacture of the pharmaceutical compositions are nontoxic.

Pharmaceutical compositions can be administered by spray inhalation. Any conventional pharmaceutical composition for spray inhalation administration may be used. Another preferred mode of administration is by aerosol.

The pharmaceutical composition of the invention can also be formulated for topical application. The composition for topical application can be in the form of an aqueous solution, lotion or jelly, an oily solution or suspension or a fatty or emulsion ointment.

5 The pharmaceutical composition of the invention can also be used for slow prolonged release with a transdermal therapeutic system in polymer matrix or with an appropriate formulation for oral slow release.

 The pharmaceutical compositions according to the present invention may also be administered orally or rectally. They may also be administered by nasal instillation,
10 aerosols or in the form of unguents or creams. The pharmaceutical compositions which can be used for oral administration may be solid or liquid, for example, in the form of uncoated or coated tablets, pills, dragees, gelatine capsules, solutions, syrups and the like. For administration by the rectal route, the compositions containing the compounds of the present invention are generally used in the form of suppositories.

15 The pharmaceutical forms, such as tablets, drops, suppositories and the like, are prepared by conventional pharmaceutical methods. The compounds of the present invention are mixed with a solid or liquid, non-toxic and pharmaceutically acceptable carrier and possibly also mixed with a dispersing agent, a disintegration agent, a stabilizing agent and the like. If appropriate, it is also possible to add preservations,
20 sweeteners, coloring agents and the like.

 Preferably, the pharmaceutical compositions of the invention is administered in traditional form for oral administration, as film coated tablets, lozenges, dragees, and oral liquid preparation such as syrup.

 Best results have been obtained with an oral dosage form, in particular liquid
25 formulations such as syrup for children, and film-coated tablet for adults. For example, patients can receive 2 doses of 0,25 mg/kg (total daily dose : 0,50 mg/kg/day) of an oral solution of levocetirizine dihydrochloride 10 mg/ml per day; one ml of the solution contains 20 drops and one drop of levocetirizine dihydrochloride solution contains 0,5 mg.

30 As an Example of a composition according to the present invention, the following formulation of a film coated tablet is preferred: levocetirizine dihydrochloride, magnesium stearate, cellulose, lactose and silicon dioxide.

 As an Example of a composition according to the present invention, the following formulation of a syrup is preferred: levocetirizine dihydrochloride, methyl-
35 and propylparaben, saccharinum, and purified water.

 Pharmaceutical compositions of the invention are useful to treat the persistent allergic rhinitis. These compositions can alleviate the effects of the persistent allergic rhinitis.

Another advantage of the invention is the ability of the process to improve quality of life and all symptoms of persistent allergic rhinitis.

The method of the invention is believed particularly suited to use in patients susceptible to suffer from persistent allergic rhinitis.

5 Another advantage of the invention is that levocetirizine dihydrochloride has an effect on rhinitis up to 6 months.

It is shown that levocetirizine dihydrochloride has an effect on quality of life up to 6 months.

10 It is shown that levocetirizine dihydrochloride has an effect on nasal congestion after 3 months. It lasts through 3 months.

The invention is further defined by reference to the following example.

Example

The aim of the study relative to the clinical effect of levocetirizine dihydrochloride was to establish on the intention to treat (ITT population) whether a 6
15 month levocetirizine dihydrochloride treatment can improve the quality of life and clinical symptoms from adult patients suffering from persistent allergic rhinitis, when compared to placebo. For clinical symptoms, it was considered that a 1 point score reduction is clinically relevant. For health-related quality of life, it was considered that a 0.36 point score reduction is relevant. Secondary parameters of efficacy included
20 different durations of treatment, different symptoms, different quality of life questionnaires, the incidence of co-morbidities suspected to be linked to allergic rhinitis and pharmaco-economic variables. The safety of this long-term treatment with levocetirizine dihydrochloride has also been evaluated.

The target population of this example consisted of adults aged more than 18
25 years suffering from persistent allergic rhinitis [WHO Initiative on Allergic Rhinitis and its Impact on Asthma (ARIA), 2000, pages S147-S149]. To be enrolled, the subjects needed to have sufficient rhinitis symptoms during the selection period. Excluded were patients with ENT or eye infection during the 2-weeks preceding initial visit.

The study was a prospective, randomized, double blind, parallel group, and
30 placebo-controlled study with levocetirizine dihydrochloride.

The severity of clinical symptoms was rated by the T5SS (sneezing, rhinorrhea, nasal pruritus, ocular pruritus and nasal congestion) evaluated each by a score from 0 to 3. The impact on health related quality of life was measured using the Rhinoconjunctivitis Quality of Life Questionnaire (RQLQ) (E. JUNIPER and G.H. GUYATT, Development and testing of a new measure of health status for clinical trials in rhinoconjunctivitis, Clinical and Experimental Allergy 1991; 21:77-83; E. JUNIPER, Measuring Health Related Quality of Life in rhinitis, J. Allergy Clin. Immunol. 1997; 99:S742-9).

Study treatment lasted for 6 months. After the treatment period, patients entered a 1-week follow-up period.

The primary end-point for efficacy was a decrease of the T5SS over the first 4 weeks of at least 1 unit of score. The primary end-point for quality of life was a
5 decrease of RQLQ after 4 weeks of at least 0.36 unit of the total score.

Secondary parameters of efficacy included the mean T5SS, the RQLQ and the SF-36 questionnaire at the different time points of the study, and the incidence and the duration of rescue medication over 6 months.

Exploratory parameters of efficacy included the mean of each individual
10 rhinitis score, each RQLQ domain and each scale of the SF-36 questionnaire at the different time points of the study, the Global Evaluation Scale after 4 weeks and 6 months, the incidence of co-morbidities suspected to be linked to allergic rhinitis and the pharmaco-economic direct and indirect costs over 6 months.

At each of the eight visits, diary book entries (T5SS, RQLQ, SF-36, indirect cost
15 pharmaco-economic parameters, concomitant medication, outpatient consultations and adverse events) were verified and transferred into the Clinical Record Form and direct cost pharmaco-economic parameters were recorded. Patients underwent a physical examination, including the measurement of vital signs. At the beginning and at the end of the study they also underwent a safety lab test, including pregnancy test
20 for females, and at Visits 4 and 7, they filled-in a Global evaluation scale.

Adverse events were recorded by the patients on diary cards and discussed with the investigator at each visit. Serious adverse events had to be reported immediately.

Oral tablets of levocetirizine dihydrochloride (5 mg) and matching placebo,
25 similar in appearance, shape and taste were used. The recommended study dosage was 1 tablet per day.

Sample size was based on 40% relative improvement over placebo in the RQLQ questionnaire, assuming an improvement from baseline for placebo of 0.9. For this questionnaire this corresponds to a difference of 0.36 vs. placebo.

30 The baseline characteristics of the two treatment groups, including demographic data, were comparable.

The study shows that treatment with levocetirizine dihydrochloride improves the symptoms of persistent allergic rhinitis (Difference of T5SS over the first 4 weeks: 1.14, $p < 0.001$; this difference being maintained over the whole study period) and the
35 QOL (Change from baseline of the RQLQ Overall Score at first 4 weeks: 0.48, $p < 0.001$; this difference being maintained over the whole study period). A statistically significant improvement is also observed at all time points for sneezing, rhinorrhea, nasal pruritus and ocular pruritus. In addition, an improvement of nasal

obstruction is observed, which becomes statistically different from 3 months onwards (Difference vs. placebo: 0.15, $p = 0.009$). Moreover, the long-term administration of levocetirizine did not involve particular safety concerns.

This study provides evidence of activity of levocetirizine in persistent allergic rhinitis. Levocetirizine is shown to be active on nasal obstruction after a long term treatment (equal or more than 3 months).

Table I

Mean T5SS over the first four weeks of treatment

(ITT population)

Treatment	N	Baseline Mean	(SD)	Mean	(SD)	Adjusted mean ^(a)	(SE)	Diff. vs. placebo ^(b) (95 % CI)	p-value ^(c)
Placebo	271	8.90	(2.26)	6.61	(2.47)	6.56	(0.15)		
Lctz 5 mg	276	9.02	(2.28)	5.53	(2.52)	5.43	(0.15)	1.14 [0.75, 1.52]	< 0.001

^(a) Mean adjusted for baseline score and country.

^(b) Placebo minus levocetirizine dihydrochloride (Lctz) 5mg.

^(c) p-value was obtained from an ANCOVA with baseline score as covariate and country and treatment as factors.

In table I it is shown that treatment with levocetirizine dihydrochloride improves the symptoms of persistent allergic rhinitis.

Table II

Change from baseline of the RQLQ Overall Score after first 4 weeks of treatment

(ITT population)

Treatment	N	Baseline		Change				p-value ^(c)
		Mean	(SD)	Mean	(SD)	Adjusted Mean ^(a)	(SE)	Diff. vs. placebo ^(b) (95 % CI)
Placebo	252	3.06	(0.94)	-0.99	(1.25)	-1.01	(0.07)	
Lctz 5 mg	257	3.04	(0.92)	-1.50	(1.18)	-1.49	(0.07)	0.48 [0.29, 0.67]

^(a) Mean adjusted for baseline score and country.

^(b) Placebo minus levocetirizine dihydrochloride 5mg.

^(c) p-value was obtained from an ANCOVA with baseline score as covariate and country and treatment as factors.

In table II it is shown that treatment with levocetirizine dihydrochloride improves the quality of life.

5

Table III

Nasal congestion symptoms evaluated over the 24 hours,
over the first week and first 4 weeks, 3, 4.5 and 6 months of treatment
(ITT population)

Period	Treatment	N	Baseline Mean (SD)	Mean (SD)	Adjusted Mean ^(a) (SE)	Diff. vs. (95 % CI) ^(b)	p- value ^(c)
Week 1	Placebo	270	1.85 (0.71)	1.64 (0.77)	1.65 (0.04)	0.07 [-0.04; 0.18]	0.203
	Lctz 5 mg	271	1.90 (0.69)	1.61 (0.83)	1.58 (0.04)		
First 4 weeks	Placebo	271	1.85 (0.71)	1.49 (0.74)	1.48 (0.04)	0.08 [-0.02; 0.19]	0.123
	Lctz 5 mg	276	1.91 (0.69)	1.44 (0.78)	1.40 (0.04)		
3 months	Placebo	270	1.85 (0.71)	1.33 (0.74)	1.31 (0.04)	0.15 [0.04; 0.26]	0.009
	Lctz 5 mg	276	1.91 (0.69)	1.22 (0.78)	1.16 (0.04)		
4.5 months	Placebo	270	1.85 (0.71)	1.29 (0.74)	1.27 (0.04)	0.15 [0.04; 0.26]	0.007
	Lctz 5 mg	276	1.91 (0.69)	1.17 (0.77)	1.11 (0.04)		
6 months	Placebo	270	1.85 (0.71)	1.26 (0.74)	1.24 (0.04)	0.16 [0.05; 0.27]	0.005
	Lctz 5 mg	276	1.91 (0.69)	1.13 (0.76)	1.08 (0.04)		

10

^(a) On study Mean adjusted for baseline score and country.

^(b) Placebo minus levocetirizine dihydrochloride.

^(c) p-value was obtained from an ANCOVA with baseline score as covariate and country and treatment as factors.

15 In table III it is shown that levocetirizine dihydrochloride is shown to be active on nasal obstruction after a long term treatment.

The following abbreviations are used in the example:

	T5SS	Total 5 Symptoms Score
	ITT	Intention-to-Treat
20	N	Number
	SD	Standard Deviation
	SE	Standard Error of the Mean
	Diff.	Difference
	vs.	versus
25	CI	Confidence Interval
	P	probability that the observed difference is only by chance
	RQLQ	Rhinoconjunctivitis Quality of Life Questionnaire
	ANCOVA	Analysis of Covariance
	ENT	Ear-Nose-Throat
30	SF-36	Medical Outcomes Survey Short Form 36
	Lctz	levocetirizine dihydrochloride.

A long duration of effect is noted. The positivity of the trial is due to the lack of tachyphylaxis, i.e there's no "adjustment" of the dosing schedule needed during 6 months. The recommended dosage is effective constantly throughout the trial.

5 An improvement of quality of life (QoL) is clearly noted during the trial. It is central to ARIA. This is the first time that a drug is able to change the QoL of patients for such a long duration. This is as close as possible to a "disease modifying" effect.

Nasal congestion is treated during the trial. Interestingly, nasal congestion is a symptom relief that appear during the trial, i.e the effect is gradual, this is congruent with the observation that QoL is improving.

10 It is demonstrated that levocetirizine dihydrochloride is able to treat persistent rhinitis as long as it is administered, but also able to modify daily activities of patients, going beyond the simple symptom relief observed in short duration trials so far.

CLAIMS

- 5 1. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the persistent allergic rhinitis.
2. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for decreasing the symptoms of
- 10 persistent allergic rhinitis and improving the quality of life.
3. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the rhinorrhea.
4. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the nasal obstruction.
- 15 5. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the nasal itching.
6. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the sneezing.
7. Use of levocetirizine or a pharmaceutically acceptable salt thereof for the preparation of a medicament intended for treating the ocular pruritis.
- 20 8. Use according to any one of claims 1 to claim 7, wherein the salt is the levocetirizine dihydrochloride.
9. Use according to any one of claims 1 to 8, wherein the medicament is adapted for administration of a daily dosage from about 0,0005 mg to about 2 mg of
- 25 said levocetirizine or said pharmaceutically acceptable salt thereof, per kg of body weight per patient.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/05491

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A61K31/495 A61P37/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, MEDLINE, BIOSIS, EMBASE, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 06429 A (SEPRACOR INC) 31 March 1994 (1994-03-31) cited in the application page 1-7 page 15 page 20 -page 21	1-9
X	<p>"Xyzal (Levocetirizine) launched in England for Allergic Rhinitis" INTERNET, 'Online! 1 October 2001 (2001-10-01), XP002240083 Retrieved from the Internet: <URL:http://www.pslgroup.com/dg/207766.htm http://www.pslgroup.com/dg/207766.htm> 'retrieved on 2003-05-05! the whole document</p> <p style="text-align: center;">-/--</p>	1-9

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

15 September 2003

Date of mailing of the international search report

23/09/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Domingues, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/05491

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>"Germany Approves Antihistamines Xyzal" INTERNET, 'Online! 16 January 2001 (2001-01-16), XP002240084 Retrieved from the Internet: <URL:http://www.pslgroup.com/dg/1efc66.htm http://www.pslgroup.com/dg/1efc66.htm> 'retrieved on 2003-05-05! the whole document</p> <p>----</p>	1-9
X	<p>GENSTHALER BM: "Levocetirizine: R-enantiomer against allergies" PHARMAZEUTISCHE ZEITUNG, vol. 146, no. 7, 15 February 2001 (2001-02-15), pages 35-36, XP001147797 page 35 page 35 -page 36</p> <p>----</p>	1-9
Y	<p>GRANT J ANDREW ET AL: "A double-blind, randomized, single-dose, crossover comparison of levocetirizine with ebastine, fexofenadine, loratadine, mizolastine, and placebo: suppression of histamine-induced wheal-and-flare response during 24 hours in healthy male subjects." ANNALS OF ALLERGY, ASTHMA & IMMUNOLOGY: OFFICIAL PUBLICATION OF THE AMERICAN COLLEGE OF ALLERGY, ASTHMA, & IMMUNOLOGY. UNITED STATES FEB 2002, vol. 88, no. 2, February 2002 (2002-02), pages 190-197, XP009010299 ISSN: 1081-1206 abstract page 195 -page 196</p> <p>----</p>	1-9
Y	<p>GANDON J M ET AL: "Lack of effect of single and repeated doses of levocetirizine, a new antihistamine drug, on cognitive and psychomotor functions in healthy volunteers." BRITISH JOURNAL OF CLINICAL PHARMACOLOGY, vol. 54, no. 1, July 2002 (2002-07), pages 51-58, XP002240086 e=bjcp July, 2002 ISSN: 0306-5251 abstract page 52 page 56 -page 58</p> <p>-----</p>	1-9

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 03/05491

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9406429	A	31-03-1994	
		AT 170749 T	15-09-1998
		AU 4932093 A	12-04-1994
		AU 711212 B2	07-10-1999
		AU 5934398 A	21-05-1998
		CA 2145410 A1	31-03-1994
		DE 69320975 D1	15-10-1998
		DE 69320975 T2	12-05-1999
		DK 663828 T3	07-06-1999
		EP 0663828 A1	26-07-1995
		EP 0950412 A2	20-10-1999
		ES 2122042 T3	16-12-1998
		JP 8501561 T	20-02-1996
		LU 90898 A9	08-04-2002
		WO 9406429 A1	31-03-1994
		US 5698558 A	16-12-1997
